

## AP<sup>®</sup> CHEMISTRY EQUATIONS AND CONSTANTS

Throughout the exam the following symbols have the definitions specified unless otherwise noted.

L, mL	= liter(s), milliliter(s)
g	= gram(s)
nm	= nanometer(s)
atm	= atmosphere(s)

mm Hg	= millimeters of mercury
J, kJ	= joule(s), kilojoule(s)
V	= volt(s)
mol	= mole(s)

### ATOMIC STRUCTURE

$$E = h\nu$$

$$c = \lambda\nu$$

$$E = \text{energy}$$

$$\nu = \text{frequency}$$

$$\lambda = \text{wavelength}$$

$$\text{Planck's constant, } h = 6.626 \times 10^{-34} \text{ J s}$$

$$\text{Speed of light, } c = 2.998 \times 10^8 \text{ m s}^{-1}$$

$$\text{Avogadro's number} = 6.022 \times 10^{23} \text{ mol}^{-1}$$

$$\text{Electron charge, } e = -1.602 \times 10^{-19} \text{ coulomb}$$

### EQUILIBRIUM

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}, \text{ where } a A + b B \rightleftharpoons c C + d D$$

$$K_p = \frac{(P_C)^c (P_D)^d}{(P_A)^a (P_B)^b}$$

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

$$K_b = \frac{[OH^-][HB^+]}{[B]}$$

$$K_w = [H^+][OH^-] = 1.0 \times 10^{-14} \text{ at } 25^\circ\text{C}$$

$$= K_a \times K_b$$

$$\text{pH} = -\log[H^+], \text{ pOH} = -\log[OH^-]$$

$$14 = \text{pH} + \text{pOH}$$

$$\text{pH} = \text{p}K_a + \log \frac{[A^-]}{[HA]}$$

$$\text{p}K_a = -\log K_a, \text{ p}K_b = -\log K_b$$

#### Equilibrium Constants

$K_c$  (molar concentrations)

$K_p$  (gas pressures)

$K_a$  (weak acid)

$K_b$  (weak base)

$K_w$  (water)

### KINETICS

$$\ln[A]_t - \ln[A]_0 = -kt$$

$$\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$$

$$t_{1/2} = \frac{0.693}{k}$$

$k$  = rate constant

$t$  = time

$t_{1/2}$  = half-life

**GO ON TO THE NEXT PAGE.**

## GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$P_A = P_{\text{total}} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{\text{total}} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = {}^{\circ}\text{C} + 273$$

$$D = \frac{m}{V}$$

$$KE \text{ per molecule} = \frac{1}{2}mv^2$$

Molarity,  $M$  = moles of solute per liter of solution

$$A = abc$$

- $P$  = pressure
- $V$  = volume
- $T$  = temperature
- $n$  = number of moles
- $m$  = mass
- $M$  = molar mass
- $D$  = density
- $KE$  = kinetic energy
- $v$  = velocity
- $A$  = absorbance
- $a$  = molar absorptivity
- $b$  = path length
- $c$  = concentration

$$\begin{aligned} \text{Gas constant, } R &= 8.314 \text{ J mol}^{-1}\text{K}^{-1} \\ &= 0.08206 \text{ L atm mol}^{-1}\text{K}^{-1} \\ &= 62.36 \text{ L torr mol}^{-1}\text{K}^{-1} \\ 1 \text{ atm} &= 760 \text{ mm Hg} = 760 \text{ torr} \\ \text{STP} &= 273.15 \text{ K and } 1.0 \text{ atm} \end{aligned}$$

## THERMODYNAMICS/ELECTROCHEMISTRY

$$q = mc\Delta T$$

$$\Delta S^\circ = \sum S^\circ \text{ products} - \sum S^\circ \text{ reactants}$$

$$\Delta H^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$= -RT \ln K$$

$$= -nFE^\circ$$

$$I = \frac{q}{t}$$

- $q$  = heat
- $m$  = mass
- $c$  = specific heat capacity
- $T$  = temperature
- $S^\circ$  = standard entropy
- $H^\circ$  = standard enthalpy
- $G^\circ$  = standard Gibbs free energy
- $n$  = number of moles
- $E^\circ$  = standard reduction potential
- $I$  = current (amperes)
- $q$  = charge (coulombs)
- $t$  = time (seconds)

Faraday's constant,  $F$  = 96,485 coulombs per mole of electrons

$$1 \text{ volt} = \frac{1 \text{ joule}}{1 \text{ coulomb}}$$